## **NEUROLOGY** ENROLLMENT FORM



Ph: 949.667.4325   Fax: 714.485.7063   altheainfusion.com						
Patient Name:		_ DOB: /	/	Height:	Weight:kg Date: / /	
Allergies:		Email:			Ship to: Patient	
Address:		Ph:	:		MD Office Prescriber Signature Althea Location:	
Prescriber:	Ph:	Fax:			Allilea Location.	
DOCUMENTS TO INCLUDE WITH	REFERRAL			NOSIS		
<ul> <li>Patient demographics</li> <li>Face sheet</li> <li>Insurance information</li> <li>Recent Lab Results (Required): An SCr, IgA level, and EMG/NCV studient H&amp;P</li> <li>Medications and therapies tried and Baseline assessment, including det</li> <li>Attach original prescription orders</li> </ul>	es I failed		De G6 G7 G6 G3 G7 G1 M3 G2 L10	rmatomyositis 1.0 Guillian-Barré Synd 0.80 Lambert-Eaton Sy 1.82 Multifocal Motor N 5 – Multiple Sclerosis (I 0.01 Myasthenia Gravis 3.0 Paraneoplastic Syn 3.22 Polymyositis 5.82 Stiff-Person Syndr 0.0 Pemphigus Vulgaris 9.3 Immune Thrombocy	ndrome europathy (MMN) RRMS subtype) s w/Acute Exacerbation drome	
IMMUNE GLOBULIN PRESCRIPTION (IVIG)		5	SUBCUTANEOUS IMMUNE GLOBULIN PRESCRIPTION (SCIG):			
grams daily  Maintenance: grams/kg i	nfused over day(s) for day(s)	F	Refill x		OR grams every weeks. select number of infusion sites and	
OK to round to the nearest vial size.+/- 4 days to allow scheduling flexibility.				doses will be admini rdered otherwise.	stered on consecutive days Non-consecutive days OK	
NEUROLOGY PRESCRIPTION	DOSAGE A	ND ADMINIST	ΓRΑΤ	ION		
Tepezza ® (teprotumumab-trbw)	10 mg/kg intravenous 8 infusions).	10 mg/kg intravenously on day 1, followed by 20 mg/kg intravenously every 3 weeks for 7 additional infusions (total of 8 infusions).				
Ocrevus® (ocrelizumab)	MAINTENANCE DOS	LOADING DOSE: Infuse Ocrevus 300mg / 250ml NS intravenously on weeks 0 and 2 (+/- 4 days). Infuse over 2.5 hours. MAINTENANCE DOSE: Infuse Ocrevus 600mg / 500ml NS intravenously over approximately 2 hours, every 6 months (+/- 7 days), starting 6 months after week 0.				
Soliris ® (eculizumab)	, ,	900 mg intravenously weekly for the first 4 weeks, followed by 1200 mg intravenously at week 5, then 1200 mg intravenously every 2 weeks thereafter.				
Rituxan® (rituximab) Ruxience® (rituximab-pvvr) Truxima® (rituximab-abbs)	1000mg intravenously	1000mg intravenously on day 1 and 15 for 2 doses, then 1000mg intravenously once every 6 to 12 months.				
Tysabri® (eculizumab)	Infuse 300mg/mcg in	Infuse 300mg/mcg intravenously every four weeks.				
Uplizna ® (inebilizumab-cdon)	300 mg intravenously	300 mg intravenously on day 1 and day 15, followed by 300 mg intravenously every 6 months thereafter.				
Vyvgart® (efgartigimod alfa-fcab)	10mg/kg intravenous	10mg/kg intravenously once weekly for 4 weeks.				
Ultomoris ® (ravulizumab-cwvz)	Administer a loading	Administer a loading dose followed 2 weeks later by weight-based maintenance dosing every 8 weeks.				
PREMEDICATION ORDERS/OTHE	R MEDICATIONS					
Flush Protocol - Flushing per S.A.S.H. p type.	rotocol (Saline, Administer n	nedication, Salin	ne, He	parin) specific volum	es and concentrations based on patient's line	
Premedications & Other Medications - In  Acetaminophen mg PO prior to		ocol. Anaphylaxis nydramine	s Kit o mg F	• •	l.	